

## New Advanced Higher Mathematics: Formulae

**Green (G):** Formulae you must memorise in order to pass Advanced Higher maths as they are not on the formula sheet.

**Amber (A):** These formulae are given on the formula sheet. But it will still be useful for you to memorise them.

**Red (R):** Don't worry about memorising these, but they might be useful to save time in classwork and homework.

### Trigonometric Identities: (from National 5 and Higher)

	<b>Essential Formulae to know off by heart for the exam (G)</b>	<b>Other useful ones that may be useful for homework/classwork etc.</b>
<b>Links between ratios</b>	$\cos^2 A + \sin^2 A = 1$ $\tan A = \frac{\sin A}{\cos A}$	$1 + \tan^2 A = \sec^2 A$ $\cot^2 A + 1 = \operatorname{cosec}^2 A$
<b>Squared</b>	$\cos^2 x = \frac{1}{2}(1 + \cos 2x)$ $\sin^2 x = \frac{1}{2}(1 - \cos 2x)$	
<b>Compound Angle</b>	$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$ $\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$	$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$
<b>Double Angle</b>	$\sin(2A) = 2 \sin A \cos A$ $\cos(2A) = \cos^2 A - \sin^2 A$	$\tan(2A) = \frac{2 \tan A}{1 - \tan^2 A}$

### Exact Values(you should know all these, though there is no non-calculator paper, unlike Higher)

	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\pi$	$\frac{3\pi}{2}$	$2\pi$	<b>Negative facts:</b> $\sin(-\theta) = -\sin(\theta)$ $\cos(-\theta) = +\cos(\theta)$ $\tan(-\theta) = -\tan(\theta)$
sin	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1	0	-1	0	
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0	-1	0	1	
tan	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	undef.	0	undef.	0	

### Complex Numbers

For the complex number,  $z = a + bi$ ,

- the **modulus** is given by  $|z| = \sqrt{a^2 + b^2}$
- and the **argument** is given by  $\tan \theta = \frac{b}{a}$   $-\pi < \theta < \pi$
- The **conjugate** is  $\bar{z} = a - bi$

**De Moivre's Theorem** says that

for any  $z = r(\cos \theta + i \sin \theta)$ , then  $z^n = r^n(\cos n\theta + i \sin n\theta)$  ( $n \in \mathbb{Q}$ )

## Differentiation

**Product Rule:**  $u \frac{dv}{dx} + v \frac{du}{dx}$       **Quotient Rule:**  $\frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$

$f(x)$	$f'(x)$	$f(x)$	$f'(x)$
$\sin^{-1} x$	$\frac{1}{\sqrt{1-x^2}}$	$\sec x$	$\sec x \tan x$
$\cos^{-1} x$	$-\frac{1}{\sqrt{1-x^2}}$	$\operatorname{cosec} x$	$-\operatorname{cosec} x \cot x$
$\tan^{-1} x$	$\frac{1}{1+x^2}$	$\cot x$	$-\operatorname{cosec}^2 x$
$\tan x$	$\sec^2 x$	$\ln f(x)$	$\frac{f'(x)}{f(x)}$
$\ln x \quad x > 0$	$\frac{1}{x}$	To differentiate an inverse function: $\frac{dx}{dy} = \frac{1}{\frac{dy}{dx}}$	
$e^x$	$e^x$		

- Parametric Equations** (where  $x = f(t), y = g(t)$ ):
- Gradient (direction of movement) =  $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}}$
  - Speed =  $\sqrt{\left(\frac{dy}{dt}\right)^2 + \left(\frac{dx}{dt}\right)^2}$
  - $\frac{d^2 y}{dx^2} = \frac{\dot{x} \ddot{y} - \dot{y} \ddot{x}}{\dot{x}^3}$

## Integration

### On Formula Sheet

$f(x)$	$\int f(x) dx$
$\sec^2 ax$	$\frac{1}{a} \tan ax + C$
$\frac{1}{\sqrt{a^2 - x^2}}$	$\sin^{-1} \left( \frac{x}{a} \right) + C$
$\frac{1}{a^2 + x^2}$	$\frac{1}{a} \tan^{-1} \left( \frac{x}{a} \right) + C$
$e^{ax}$	$\frac{1}{a} e^{ax} + C$

To save you time in hard questions for homework/classwork, **no need to memorise:**

$f(x)$	$\int f(x) dx$
$\tan x$	$\ln  \sec x  + C$
$\operatorname{cosec} x$	$-\ln  \operatorname{cosec} x + \cot x  + C$
$\cot x$	$\ln  \sin x  + C$
$\sec x$	$\ln  \sec x + \tan x  + C$

### Integration by Parts

$$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$$

Volume of solid of revolution  $f(x)$  between  $a$  and  $b$ :

About x axis:  $V = \pi \int_a^b f(x)^2 dx$       About y axis:  $V = \pi \int_a^b f(y)^2 dy$

**Sequences and Series**

	Arithmetic Series	Geometric Series
<b><math>n^{\text{th}}</math> term</b>	$u_n = a + (n-1)d$	$u_n = ar^{n-1}$
<b>Sum of <math>n</math> terms</b>	$S_n = \frac{1}{2}n(2a + (n-1)d)$	$S_n = \frac{a(1-r^n)}{1-r} \quad r \neq 1$
<b>Sum to infinity</b>		$S_\infty = \frac{a}{1-r} \quad  r  < 1$

**Important Identities**

$$\sum_{k=1}^n 1 = n$$

$$\sum_{r=1}^n r = \frac{n(n+1)}{2} \qquad \sum_{r=1}^n r^2 = \frac{n(n+1)(2n+1)}{6}$$

$$\sum_{r=1}^n r^3 = \frac{n^2(n+1)^2}{4}$$

**Maclaurin Series**

$$f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \dots + \frac{f^{(n)}(0)}{n!}x^n + \dots$$

and in particular:

**Very useful to memorise:**

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!} + \dots$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$$

**Less essential to memorise:**

$$\tan^{-1} x = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots$$

$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$

**Functions**

**Odd function:**  $f(-x) = -f(x)$   
(180° rotational symmetry)

**Even function:**  $f(-x) = f(x)$   
(line symmetry about the  $y$ -axis)

**Binomial Theorem**

The coefficient of the  $r^{\text{th}}$  term in the binomial expansion  $(x + y)^n$  is  $\binom{n}{r} x^{n-r} y^r$

$${}^n C_r = \binom{n}{r} = \frac{n!}{r!(n-r)!}$$

**Vectors, Lines and Planes**

Angle between two vectors: (Higher)  $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos \theta$

**Equations of a 3d line:** through  $(x_1, y_1, z_1)$  and with direction vector  $\mathbf{d} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$

Parametric form

$$\begin{aligned} x &= x_1 + at \\ y &= y_1 + bt \quad (\mathbf{x} = \mathbf{a} + t\mathbf{d}) \\ z &= z_1 + ct \end{aligned}$$

Symmetric form

$$\frac{x - x_1}{a} = \frac{y - y_1}{b} = \frac{z - z_1}{c} (= t)$$

**Equations of a plane:**

Normal  $\mathbf{n}$  is  $\begin{pmatrix} l \\ m \\ n \end{pmatrix}$

Point on line = P (with position vector  $\mathbf{a}$ )

Vector equation

$$\mathbf{x} \cdot \mathbf{n} = \mathbf{a} \cdot \mathbf{n}$$

Symmetric/Cartesian

$$lx + my + nz = k$$

where  $k = \mathbf{a} \cdot \mathbf{n}$

Parametric (A)

$$\mathbf{x} = \mathbf{a} + s\mathbf{b} + t\mathbf{c}$$

( $\mathbf{b}$  and  $\mathbf{c}$  are any two non-parallel vectors in plane)

**Angle between two lines** = Acute angle between their direction vectors

**Angle between two planes** = Acute angle between their normals

**Angle between line and plane** =  $90^\circ -$  (Acute angle between  $\mathbf{n}$  and  $\mathbf{d}$ )

**Cross (vector) product:**

$$\mathbf{a} \times \mathbf{b} = |\mathbf{a}||\mathbf{b}| \sin \theta \hat{\mathbf{n}} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = \mathbf{i} \begin{vmatrix} a_2 & a_3 \\ b_2 & b_3 \end{vmatrix} - \mathbf{j} \begin{vmatrix} a_1 & a_3 \\ b_1 & b_3 \end{vmatrix} + \mathbf{k} \begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix}$$

**Scalar triple product:**

$$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

## Matrices

		Determinant and Inverse
<b>2×2 matrices</b>	$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$	$\det A = ad - bc$ and $A^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$
<b>3×3 matrices</b>	$A = \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix}$	$\det A = a \begin{vmatrix} e & f \\ h & i \end{vmatrix} - b \begin{vmatrix} d & f \\ g & i \end{vmatrix} + c \begin{vmatrix} d & e \\ g & h \end{vmatrix}$

$$(AB)^{-1} = B^{-1}A^{-1} \qquad (AB)^T = B^T A^T \qquad \det AB = \det A \det B \quad (A)$$

### Transformation Matrices

Anti-CW Rotation by $\theta$ degrees	$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$ ,	Reflection in $y$ -axis	$\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$
Dilatation by scale factor $a$	$\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$ ,	Reflection in $x$ -axis	$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$

## Differential Equations

For  $\frac{dy}{dx} + P(x)y = Q(x)$ , the Integrating Factor  $I(x)$  is  $e^{\int P(x)dx}$

and the solution is given by  $I(x)y = \int I(x)Q(x)dx$

### Second Order Differential Equations

#### COMPLEMENTARY FUNCTION (Homogeneous Equations)

Nature of roots	Form of general solution
Two distinct real $m$ and $n$	$y = Ae^{mx} + Be^{nx}$
Real and equal $m$	$y = (A + Bx)e^{mx}$
Complex conjugate $m = p \pm iq$	$y = e^{px} (A \cos qx + B \sin qx)$

#### PARTICULAR INTEGRAL (Inhomogeneous Equations)

Right-hand side contains...	For Particular Integral, try...
$\sin ax$ or $\cos ax$	$y = P \cos ax + Q \sin ax$
$e^{ax}$	$y = Pe^{ax}$
Linear expression $y = ax + b$	$y = Px + Q$
Quadratic expression $y = ax^2 + bx + c$	$y = Px^2 + Qx + R$