

2006 Mathematics

Higher – Paper 1

Finalised Marking Instructions

© The Scottish Qualifications Authority 2006

The information in this publication may be reproduced to support SQA qualifications only on a non-commercial basis. If it is to be used for any other purposes written permission must be obtained from the Assessment Materials Team, Dalkeith.

Where the publication includes materials from sources other than SQA (secondary copyright), this material should only be reproduced for the purposes of examination or assessment. If it needs to be reproduced for any other purpose it is the centre's responsibility to obtain the necessary copyright clearance. SQA's Assessment Materials Team at Dalkeith may be able to direct you to the secondary sources.

These Marking Instructions have been prepared by Examination Teams for use by SQA Appointed Markers when marking External Course Assessments. This publication must not be reproduced for commercial or trade purposes.

1. Marks must be assigned in accordance with these marking instructions. In principle, marks are awarded for what is correct, rather than marks deducted for what is wrong.
2. Award one mark for each ‘bullet’ point. Each error should be underlined in RED at the point in the working where it first occurs, and not at any subsequent stage of the working.
3. The working subsequent to an error must be followed through by the marker with possible full marks for the subsequent working, provided that the difficulty involved is approximately similar. Where, subsequent to an error, the working is eased, a deduction(s) of mark(s) should be made.
This may happen where a question is divided into parts. In fact, failure to even answer an earlier section does not preclude a candidate from assuming the result of that section and obtaining full marks for a later section.
4. Correct working should be ticked (\checkmark). This is essential for later stages of the SQA procedures. Where working subsequent to an error(s) is correct and scores marks, it should be marked with a crossed tick (\times or $X\checkmark$). In appropriate cases attention may be directed to work which is not quite correct (e.g. bad form) but which has not been penalised, by underlining with a dotted or wavy line.
Work which is correct but inadequate to score any marks should be corrected with a double cross tick ($\times\times$).
5.
 - The total mark for each section of a question should be entered in red in the **outer** right hand margin, opposite the end of the working concerned.
 - Only the mark should be written, **not** a fraction of the possible marks.
 - These marks should correspond to those on the question paper and these instructions.
6. It is of great importance that the utmost care should be exercised in adding up the marks. Where appropriate, all summations for totals and grand totals must be carefully checked. Where a candidate has scored zero marks for any question attempted, “0” should be shown against the answer.
7. As indicated on the front of the question paper, full credit should only be given where the solution contains appropriate working. Accept answers arrived at by inspection or mentally where it is possible for the answer so to have been obtained. Situations where you may accept such working will normally be indicated in the marking instructions.
8. Do not penalise:

• working subsequent to a correct answer	• omission of units
• legitimate variations in numerical answers	• bad form
• correct working in the “wrong” part of a question	

9. No piece of work should be scored through without careful checking - even where a fundamental misunderstanding is apparent early in the answer. Reference should always be made to the marking scheme - answers which are widely off-beam are unlikely to include anything of relevance but in the vast majority of cases candidates still have the opportunity of gaining the odd mark or two provided it satisfies the criteria for the mark(s).
10. If in doubt between two marks, give an intermediate mark, but without fractions. When in doubt between consecutive numbers, give the higher mark.
11. In cases of difficulty covered neither in detail nor in principle in the Instructions, attention may be directed to the assessment of particular answers by making a referral to the P.A. Please see the general instructions for P.A. referrals.
12. No marks should be deducted at this stage for careless or badly arranged work. In cases where the writing or arrangement is very bad, a note may be made on the upper left-hand corner of the front cover of the script.
13. Transcription errors: In general, as a consequence of a transcription error, candidates lose the opportunity of gaining either the first ic mark or the first pr mark.
14. Casual errors: In general, as a consequence of a casual error, candidates lose the opportunity of gaining the appropriate ic mark or pr mark.
15. **Do not write any comments on the scripts.** A revised summary of acceptable notation is given on page 4.
16. Working that has been crossed out by the candidate cannot receive any credit. If you feel that a candidate has been disadvantaged by this action, make a P.A. Referral.
17. Throughout this paper, unless specifically mentioned, a correct answer with no working receives no credit.

Summary

Throughout the examination procedures many scripts are remarked. It is essential that markers follow common procedures:

- 1 **Tick** correct working.
- 2 Put a mark in the **outer right-hand margin to match the marks allocations on the question paper.**
- 3 Do **not** write marks as fractions.
- 4 Put each mark **at the end** of the candidate's response to the question.
- 5 **Follow through** errors to see if candidates can score marks subsequent to the error.
- 6 Do **not** write any comments on the scripts.

Higher Mathematics : A Guide to Standard Signs and Abbreviations

Remember - No comments on the scripts. Please use the following and nothing else.


Signs

- ✓ The tick. You are not expected to tick every line but of course you must check through the whole of a response.


- ✕ The cross and underline. Underline an error and place a cross at the end of the line.

- ✕ The tick-cross. Use this to show correct work where you are **following through** subsequent to an error.

- ∧ The roof. Use this to show something is missing such as a crucial step in a proof or a 'condition' etc.

-  The tilde. Use this to indicate a minor transgression which is not being penalised (such as bad form).

- ✕ The double cross-tick. Use this to show correct work but which is inadequate to score any marks. This may happen when working has been eased.

		margins	
$\frac{dy}{dx} = 4x - 7$ $4x - 7 = 0$ $x = \frac{7}{4}$ $y = 3\frac{7}{8}$	✓ • ✕ ✕ •		2
$C = (1, -1)$ $m = \frac{3 - (-1)}{4 - 1}$ $m_{rad} = \frac{4}{3}$ $m_{tgt} = \frac{-1}{\frac{4}{3}}$ $m_{tgt} = -\frac{3}{4}$ $y - 3 = -\frac{3}{4}(x - 2)$	✕ ✕ • ✕ • ✕ •		3
$x^2 - 3x = 28$ $x = 7$	✓ • ∧ ✕		1
$\sin(x) = 0.75 = inv \sin(0.75) = 48.6^\circ$	 ✓ •		1

Remember - No comments on the scripts. No abbreviations. No new signs. Please use the above and nothing else.


All of these are to help us be more consistent and **accurate**.

Note: There is no such thing as a transcription error, a trivial error, a casual error or an insignificant error. These are all mistakes and as a consequence a mark is lost.

Page 5 lists the syllabus coding for each topic. This information is given in the legend underneath the question. The calculator classification is CN(calculator neutral), CR(calculator required) and NC(non-calculator).

1	2	UNIT 1	1	2	UNIT 2	1	2	UNIT 3	Year
		A1	determine range/domain		A15	use the general equation of a parabola		A28	use the laws of logs to simplify/find equiv. expression
		A2	recognise general features of graphs:poly,exp,log		A16	solve a quadratic inequality		A29	sketch associated graphs
		A3	sketch and annotate related functions		A17	find nature of roots of a quadratic		A30	solve equs of the form $A = Be^{kt}$ for A,B,k or t
		A4	obtain a formula for composite function		A18	given nature of roots, find a condition on coeffs		A31	solve equs of the form $\log_b(a) = c$ for a,b or c
		A5	complete the square		A19	form an equation with given roots		A32	solve equations involving logarithms
		A6	interpret equations and expressions		A20	apply A15-A19 to solve problems		A33	use relationships of the form $y = ax^n$ or $y = ab^x$
		A7	determine function(poly,exp,log) from graph & vv					A34	apply A28-A33 to problems
		A8	sketch/annotate graph given critical features						
		A9	interpret loci such as st.lines,para,poly, circle						
		A10	use the notation u_n for the nth term		A21	use Rem Th. For values, factors, roots		G16	calculate the length of a vector
		A11	evaluate successive terms of a RR		A22	solve cubic and quartic equations		G17	calculate the 3rd given two from A,B and vector AB
		A12	decide when RR has limit/interpret limit		A23	find intersection of line and polynomial		G18	use unit vectors
		A13	evaluate limit		A24	find if line is tangent to polynomial		G19	use: if \mathbf{u}, \mathbf{v} are parallel then $\mathbf{v} = k\mathbf{u}$
		A14	apply A10-A14 to problems		A25	find intersection of two polynomials		G20	add, subtract, find scalar mult. of vectors
					A26	confirm and improve on approx roots		G21	simplify vector pathways
					A27	apply A21-A26 to problems		G22	interpret 2D sketches of 3D situations
								G23	find if 3 points in space are collinear
								G24	find ratio which one point divides two others
		G1	use the distance formula		G9	find C/R of a circle from its equation/other data		G25	given a ratio, find/interpret 3rd point/vector
		G2	find gradient from 2 pts./angle/equ. of line		G10	find the equation of a circle		G26	calculate the scalar product
		G3	find equation of a line		G11	find equation of a tangent to a circle		G27	use: if \mathbf{u}, \mathbf{v} are perpendicular then $\mathbf{v} \cdot \mathbf{u} = 0$
		G4	interpret all equations of a line		G12	find intersection of line & circle		G28	calculate the angle between two vectors
		G5	use property of perpendicular lines		G13	find if/when line is tangent to circle		G29	use the distributive law
		G6	calculate mid-point		G14	find if two circles touch		G30	apply G16-G29 to problems eg geometry probs.
		G7	find equation of median, altitude, perp. bisector		G15	apply G9-G14 to problems			
		G8	apply G1-G7 to problems eg intersect.,concur.,collim.						
		C1	differentiate sums, differences		C12	find integrals of px^n and sums/diffs		C20	differentiate $p\sin(ax+b)$, $p\cos(ax+b)$
		C2	differentiate negative & fractional powers		C13	integrate with negative & fractional powers		C21	differentiate using the chain rule
		C3	express in differentiable form and differentiate		C14	express in integrable form and integrate		C22	integrate $(ax + b)^n$
		C4	find gradient at point on curve & vv		C15	evaluate definite integrals		C23	integrate $p\sin(ax+b)$, $p\cos(ax+b)$
		C5	find equation of tangent to a polynomial/trig curve		C16	find area between curve and x-axis		C24	apply C20-C23 to problems
		C6	find rate of change		C17	find area between two curves			
		C7	find when curve strictly increasing etc		C18	solve differential equations(variables separable)			
		C8	find stationary points/values		C19	apply C12-C18 to problems			
		C9	determinenature of stationary points						
		C10	sketch curvegiven the equation						
		C11	apply C1-C10 to problems eg optimise, greatest/least						
		T1	use gen. features of graphs of $f(x)=k\sin(ax+b)$, $f(x)=k\cos(ax+b)$; identify period/amplitude		T7	solve linear & quadratic equations in radians		T12	solve sim.equs of form $k\cos(a)=p$, $k\sin(a)=q$
		T2	use radians inc conversion from degrees & vv		T8	apply compound and double angle (c & da) formulae in numerical & literal cases		T13	express $p\cos(x)+q\sin(x)$ in form $k\cos(x\pm a)$ etc
		T3	know and use exact values		T9	apply c & da formulae in geometrical cases		T14	find max/min/zeros of $p\cos(x)+q\sin(x)$
		T4	recognise form of trig. function from graph		T10	use c & da formulaewhen solving equations		T15	sketch graph of $y=p\cos(x)+q\sin(x)$
		T5	interpret trig. equations and expressions		T11	apply T7-T10 to problems		T16	solve equ of the form $y=p\cos(rx)+q\sin(rx)$
		T6	apply T1-T5 to problems					T17	apply T12-T16 to problems

- 1 Triangle ABC has vertices A(-1,12), B(-2, -5) and C(7, -2).
- (a) Find the equation of the median BD.
- (b) Find the equation of the altitude AE.
- (c) Find the coordinates of the point of intersection of BD and AE.



3
3
3

Qu.	part	marks	Grade	Syllabus Code	Calculator class	Source
1	a,b,c	3,3,3	C	G7, G8	CN	06/01

The primary method m/s is based on the following generic m/s. THIS GENERIC M/S MAY BE USED AS AN EQUIVALENCE GUIDE BUT ONLY WHERE A CANDIDATE DOES NOT USE THE PRIMARY METHOD OR ANY ALTERNATIVE METHOD SHOWN IN DETAIL IN THE MARKING SCHEME

- ¹ ic interpret “median”
- ² ss find gradient
- ³ ic state equation
- ⁴ ss find gradient
- ⁵ ss find perpendicular gradient
- ⁶ ic state equation
- ⁷ ss start to solve simultaneous equations
- ⁸ pr solve for one variable
- ⁹ pr process

Primary Method : Give 1 mark for each •

- ¹ $D = (3,5)$
- ² $m_{BD} = 2$
- ³ $y - 5 = 2(x - 3)$ or $y + 5 = 2(x - (-2))$ etc **3 marks**
- ⁴ $m_{BC} = \frac{1}{3}$ **stated explicitly**
- ⁵ $m_{alt} = -3$
- ⁶ $y - 12 = -3(x - (-1))$ **3 marks**
- ⁷ $y - 5 = 2(x - 3)$ **and** $y - 12 = -3(x - (-1))$ **or equivalent**
- ⁸ $x = 2$
- ⁹ $y = 3$ **3 marks**

Notes

- 1 For candidates who find two medians •¹, •², •³ and •⁷, •⁸, •⁹ are available.
- 2 For candidates who find two altitudes •⁴, •⁵, •⁶ and •⁷, •⁸, •⁹ are available.
- 3 For candidates who find (a) altitude and (b) median see common error box number 3.
- 4 In (a) note that (4, 7) happens to lie on the median but does not qualify as a point to be used in •³.

Notes cont

- 5 In (b) •⁶ is only available as a consequence of attempting to find a perpendicular gradient.
- 6 In (b) candidates who guess the coordinates for E and use these to find the equation AE, can earn no marks in this part.
- 7 In (c) note that “equating zeros” is only a valid strategy when either the coefficients of x or the coefficients of y are equal.
- 8 •⁷ is a strategy mark for juxtaposing the two required equations.
- 9 See general note at the foot of page 7.

**Common Error 1
Finding two medians**

- ¹ $D = (3,5)$
 - ² $m_{BD} = 2$
 - ³ $y - 5 = 2(x - 3)$
 - ⁴ X
 - ⁵ X
 - ⁶ X
 - ⁷ $y = 2x - 1$ & $31x + 7y = 53$
 - ⁸ $x = \frac{4}{3}$
 - ⁹ $y = \frac{5}{3}$
- maximum of 6 marks

**Common Error 2
Finding two altitudes**

- ¹ X
 - ² X
 - ³ X
 - ⁴ $m_{BC} = \frac{1}{3}$
 - ⁵ $m_{alt} = -3$
 - ⁶ $y - 12 = -3(x - (-1))$
 - ⁷ $4x - 7y = 27$ & $y = -3x + 9$
 - ⁸ $x = \frac{18}{5}$
 - ⁹ $y = -\frac{9}{5}$
- maximum of 6 marks


**Common Error 3
Finding (a) altitude and (b) median**

- ¹ $m_{AC} = -\frac{7}{4}$
 - X ✓ •² $m_{BD} = \frac{4}{7}$
 - ³ $y - 5 = \frac{4}{7}(x - -2)$
 - X ✓ •⁴ *midpt of BC* = $(\frac{5}{2}, -\frac{7}{2})$
 - ⁵ $m_{AC} = -\frac{31}{7}$
 - ⁶ $y - 12 = -\frac{31}{7}(x - (-1))$
 - X ✓ •⁷ $4x - 7y = 27$ & $31x + 7y = 53$
 - X ✓ •⁸ $x = \frac{16}{7}$
 - X ✓ •⁹ $y = -\frac{125}{49}$
- maximum of 5 marks

2 A circle has centre $C(-2, 3)$ and passes through $P(1, 6)$.

(a) Find the equation of the circle.

(b) PQ is a diameter of the circle. Find the equation of the tangent to this circle at Q .



2
4

Qu.	part	marks	Grade	Syllabus Code	Calculator class	Source
2	a	2	C	G10	CN	06/54
	b	4	C	G11	CN	

The primary method m/s is based on the following generic m/s. THIS GENERIC M/S MAY BE USED AS AN EQUIVALENCE GUIDE BUT ONLY WHERE A CANDIDATE DOES NOT USE THE PRIMARY METHOD OR ANY ALTERNATIVE METHOD SHOWN IN DETAIL IN THE MARKING SCHEME

- ¹ ic enter coord. of centre in general equation
- ² ss find $(\text{radius})^2$
- ³ ss e.g. use $\overrightarrow{PC} = \overrightarrow{CQ}$ to find Q
- ⁴ pr find gradient of diameter
- ⁵ ss know and use tangent perp. to diameter
- ⁶ ic state equation

Primary Method : Give 1 mark for each •

- $(x - a)^2 + (y - b)^2 = r^2$
- ¹ $(x - (-2))^2 + (y - 3)^2$
- ² $r^2 = 18$ 2 marks
- ³ $Q = (-5, 0)$
- ⁴ $m_{\text{diameter}} = 1$ stated or implied by •5
- ⁵ $m_{\text{tangent}} = -1$
- ⁶ $y - 0 = -(x - (-5))$ 4 marks

Notes

- 1 In (a) $(\sqrt{18})^2$ is not acceptable for •².
- 2 In (b) if the coordinates of Q are estimated (i.e. guessed) then •⁶ can only be awarded if the coordinates are of the form $(a, 0)$ where $a < -2$.
- 3 In (b) •⁶ is only available if an attempt has been made to find a perpendicular gradient.

Alternative Method for (a)

- For answers of the form $x^2 + y^2 + 2gx + 2fy + c = 0$
- ¹ $x^2 + y^2 + 4x - 6y + c = 0$
 - ² $c = -5$

General Notes applicable throughout the marking scheme

There are many instances when follow throughs come into play and these will not always be highlighted for you. The following example is a reminder of what you have to look out for when you are marking.

example

At the •³ stage a candidate start with the wrong coordinates for Q . Then

$$\begin{array}{l}
 X \quad \bullet^3 \quad Q = (-4, 0) \\
 X \checkmark \quad \bullet^4 \quad m_{\text{diameter}} = \frac{6}{5} \\
 X \checkmark \quad \bullet^5 \quad m_{\text{tangent}} = -\frac{5}{6} \\
 X \checkmark \quad \bullet^6 \quad y - 0 = -\frac{5}{6}(x - (-4))
 \end{array}$$

so the candidate loses •³ but gains •⁴, •⁵ and •⁶ as a consequence of following through.

Any error can be followed through and the subsequent marks awarded provided the working has not been eased.

Any deviation from this will be noted in the marking scheme.

3	Two functions f and g are defined on the set of real numbers by $f(x) = 2x + 3$ and $g(x) = 2x - 3$.		
(a)	Find an expressions for (i) $f(g(x))$ (ii) $g(f(x))$.		3
(b)	Determine the least possible value of $f(g(x)) \times g(f(x))$.		2

Qu.	part	marks	Grade	Syllabus Code	Calculator class	Source
3	a	3	C	A4	CN	06/07
	b	2	C	A6	CN	

The primary method m/s is based on the following generic m/s. THIS GENERIC M/S MAY BE USED AS AN EQUIVALENCE GUIDE BUT ONLY WHERE A CANDIDATE DOES NOT USE THE PRIMARY METHOD OR ANY ALTERNATIVE METHOD SHOWN IN DETAIL IN THE MARKING SCHEME

- ¹ ic int. composition
- ² ic int. composition
- ³ ic int. composition
- ⁴ pr simplify all functions
- ⁵ ic int. result

Primary Method : Give 1 mark for each •

- ¹ $f(g(x)) = f(2x - 3)$ **stated or implied by •2**
- ² $2(2x - 3) + 3$
- ³ $g(f(x)) = 2(2x + 3) - 3$ **3 marks**
- ⁴ $16x^2 - 9$ **stated explicitly**
- ⁵ min.value = -9 **2 marks**

Notes

- 1 In (a) 2 marks are available for finding one of $f(g(x))$ or $g(f(x))$ and the third mark is for the other one.
- 2 In (a) the finding of $f(f(x))$ and $g(g(x))$ earns no marks.
- 3 •⁵ is only available if •⁴ has been awarded.
- 4 In (b) for •⁵, no justification is necessary. Ignore any comments, rational or irrational.

Alternative Marking 1 [Marks 1-3]

- ¹ $g(f(x)) = g(2x + 3)$
- ² $2(2x + 3) - 3$
- ³ $f(g(x)) = 2(2x - 3) + 3$

Common Error No.1 for (a) "g and f" transposed.

- | | | |
|-----|----------------|---------------------------|
| X | • ¹ | $f(g(x)) = f(2x + 3)$ |
| ✓ X | • ² | $2(2x + 3) - 3$ |
| ✓ X | • ³ | $g(f(x)) = 2(2x - 3) + 3$ |
- Award 2 out of 3

Common Error No.2 for (a)

- | | | |
|-----|----------------|---------------------------|
| X | • ¹ | $f(g(x)) = f(2x + 3)$ |
| ✓ X | • ² | $2(2x + 3) - 3$ |
| ✓ | • ³ | $g(f(x)) = 2(2x + 3) - 3$ |
- Award 2 out of 3

Common Error No.3 for (a) Repeated error

- | | | |
|-----|----------------|---------------------------|
| ✓ | • ¹ | $f(g(x)) = f(2x - 3)$ |
| X | • ² | $2(2x + 3) - 3$ |
| ✓ X | • ³ | $g(f(x)) = 2(2x - 3) + 3$ |
- Award 2 out of 3

4 A sequence is defined by the recurrence relation $u_{n+1} = 0.8u_n + 12$, $u_0 = 4$.

(a) State why the recurrence relation has a limit. 1

(b) Find this limit. 2

Qu.	part	marks	Grade	Syllabus Code	Calculator class	Source
4	a	1	C	A12	NC	06/28
	b	2	C	A13	NC	

The primary method m/s is based on the following generic m/s. THIS GENERIC M/S MAY BE USED AS AN EQUIVALENCE GUIDE BUT ONLY WHERE A CANDIDATE DOES NOT USE THE PRIMARY METHOD OR ANY ALTERNATIVE METHOD SHOWN IN DETAIL IN THE MARKING SCHEME

- ¹ ic state limit condition
- ² ss know how to find L
- ³ pr process limit

Primary Method : Give 1 mark for each •

- ¹ sequence has limit since $-1 < 0.8 < 1$ 1 mark
- ² $L = 0.8L + 12$
- ³ limit = 60 2 marks

Notes

For (a)

1 **Accept**

$$|0.8| < 1$$

$$0 < 0.8 < 1$$

0.8 lies between -1 and 1

0.8 is a proper fraction

2 **Do NOT accept**

$$-1 \leq 0.8 \leq 1$$

$-1 < a < 1$ unless a is clearly identified/replaced by 0.8 anywhere in the answer.

$$0.8 < 1$$

In (b)

3 $L = \frac{b}{1-a}$ and nothing else gains **no** marks.

4 $L = \frac{12}{0.2}$ or $\frac{120}{2}$ or $\frac{60}{1}$ etc does **NOT** gain •³.

5 An answer of 60 without any working gains **NO** marks.

6 Any calculations based on "wrong" formulae gain **NO** marks.

Alternative Method for (b)

$$\bullet^2 L = \frac{12}{1-0.8}$$

$$\bullet^3 \text{ limit} = 60$$

Bad Form

$$\bullet^2 L = \frac{12}{0.2}$$

$$\bullet^3 \text{ limit} = 60$$

award 2 marks

Common Error 1

$$X \bullet^2 L = \frac{4}{1-0.8}$$

$$X \checkmark \bullet^3 \text{ limit} = 20$$

- 5 A function f is defined by $f(x) = (2x - 1)^5$. Find the coordinates of the stationary point on the graph with equation $y = f(x)$ and determine its nature.

7

Qu.	part	marks	Grade	Syllabus Code	Calculator class	Source
5		6	C	C8, C9	NC	06/76
		1	B			

The primary method m/s is based on the following generic m/s. THIS GENERIC M/S MAY BE USED AS AN EQUIVALENCE GUIDE BUT ONLY WHERE A CANDIDATE DOES NOT USE THE PRIMARY METHOD OR ANY ALTERNATIVE METHOD SHOWN IN DETAIL IN THE MARKING SCHEME

- ¹ ss know to start to differentiate
- ² pr differentiate
- ³ ss set derivative = 0
- ⁴ pr solve
- ⁵ pr evaluate
- ⁶ ic justification
- ⁷ ic state conclusion

Primary Method : Give 1 mark for each •

- ¹ $f'(x) = \dots\dots$
- ² $5(2x - 1)^4 \times 2$
- ³ $f'(x) = 0$
- ⁴ $x = \frac{1}{2}$
- ⁵ $f(\frac{1}{2}) = 0$
- ⁶ nature table
- ⁷ pt of inflexion at $(\frac{1}{2}, 0)$

7 marks

Notes

- 1 The “= 0” shown at •³ must appear at least once somewhere in the working between •¹ and •⁴ (but not necessarily at •³).
- 2 •⁴ is only available as a consequence of solving $f'(x) = 0$.
- 3 A wrong derivative which eases the working will preclude at least •⁴ from being awarded.
- 4 For marks •⁶ and •⁷, a nature table is mandatory. The minimum amount of detail that is required is shown here:

	$< \frac{1}{2}$	$\frac{1}{2}$	$> \frac{1}{2}$
$f'(x)$	+	0	+
	∴	∴	∴

Candidates who use only $f''(x) = 0$ and try to draw conclusions from this cannot gain •⁶ or •⁷.

[$f''(x) = 0$ is a necessary but not sufficient condition for identifying points of inflexion].

- 5 •⁷ is **ONLY** available subsequent to a correct nature table for the candidate's own derivative.
- 6 •⁴ is lost in each of the following cases for the candidate's solution to the equation at •³.
 - (i) $x = \frac{1}{2}$ and $x = \text{something else}$
 - (ii) two wrong values for x
 - (iii) guess a value for x

Only one value for x needs to be followed through for •⁵, •⁶ and •⁷.


Common Error No.1

- | | | |
|--|----------------|----------------------|
| ✓ | • ¹ | $f'(x) = \dots\dots$ |
| X | • ² | $5(2x - 1)^4$ |
| ✓ | • ³ | $f'(x) = 0$ |
| X ✓ | • ⁴ | $x = \frac{1}{2}$ |
| • ⁵ , • ⁶ and • ⁷ are still available | | |

Common Error No.2

- | | | |
|--|----------------|--------------------------|
| ✓ | • ¹ | $f'(x) = \dots\dots$ |
| X | • ² | $\frac{1}{12}(2x - 1)^6$ |
| ✓ | • ³ | $f'(x) = 0$ |
| X ✓ | • ⁴ | $x = \frac{1}{2}$ |
| • ⁵ , • ⁶ and • ⁷ are still available | | |

- 6 The graph shown has equation $y = x^3 - 6x^2 + 4x + 1$.
 The shaded area is bounded by the curve, the x -axis,
 the y -axis and the line $x = 2$.
- (a) Calculate the shaded area labelled S.
 (b) Hence find the total shaded area.



4
3

Qu.	part	marks	Grade	Syllabus Code	Calculator class	Source
6	a	4	C	C16	NC	06/40
	b	3	B	C16	NC	

The primary method m/s is based on the following generic m/s. THIS GENERIC M/S MAY BE USED AS AN EQUIVALENCE GUIDE BUT ONLY WHERE A CANDIDATE DOES NOT USE THE PRIMARY METHOD OR ANY ALTERNATIVE METHOD SHOWN IN DETAIL IN THE MARKING SCHEME

- ¹ ss know to integrate
- ² pr integrate
- ³ ic substitute limits
- ⁴ pr evaluate
- ⁵ ic use result from •² with new limits
- ⁶ pr evaluate
- ⁷ ss deal with the “-ve” sign and evaluate total area

Primary Method : Give 1 mark for each •

- ¹ $\int_0^1 (x^3 - 6x^2 + 4x + 1) dx$ stated or implied by •²
- ² $\frac{1}{4}x^4 - \frac{6}{3}x^3 + \frac{4}{2}x^2 + x$
- ³ $\left(\frac{1}{4} \cdot 1^4 - 2 \cdot 1^3 + 2 \cdot 1^2 + 1\right) - 0$
- ⁴ $\frac{5}{4}$ or equivalent 4
- ⁵ $\int_1^2 \dots dx$
- ⁶ $\left(\frac{1}{4} \cdot 2^4 - 2 \cdot 2^3 + 2 \cdot 2^2 + 2\right) - \left(\frac{1}{4} \cdot 1^4 - 2 \cdot 1^3 + 2 \cdot 1^2 + 1\right) = -\frac{13}{4}$
- ⁷ $\frac{9}{2}$ or equivalent 3

Notes

for (a)

- 1 Only a limited number of marks are available to candidates who differentiate –see Common Error No.1.
- 2 In (a) candidates who transpose the limits can still earn •⁴ if the deal with the “-ve” sign appropriately.
- 3 In (b)
 - ⁷ is lost for such statements as $-3\frac{1}{4} = 3\frac{1}{4}$.

- 4 In (b) using $\int_0^2 \dots dx$ earns no marks.

Common Error No.1

- ✓ •¹ $\int_0^1 (x^3 - 6x^2 + 4x + 1) dx$
- X •² $3x^2 - 12x + 4$
- X •³ $(3 \cdot 1^2 - 12 \cdot 1 + 4) - 4$
- X •⁴ -9
- ✓ •⁵ $\int_1^2 \dots dx$ or equivalent
- X ✓ •⁶ $(3 \cdot 2^2 - 12 \cdot 2 + 4) - (3 \cdot 1^2 - 12 \cdot 1 + 4) = -3$
- X ✓ •⁷ 12

Alternative Method 1 for (b)

- ⁵ $\int_2^1 \dots dx$
- ⁶ $\left(\frac{1}{4} \cdot 1^4 - 2 \cdot 1^3 + 2 \cdot 1^2 + 1\right) - \left(\frac{1}{4} \cdot 2^4 - 2 \cdot 2^3 + 2 \cdot 2^2 + 2\right)$
- ⁷ $\frac{9}{2}$

Alternative Method 2 for (b)

- ⁵ $-\int_1^2 \dots dx$
- ⁶ $-\left(\frac{1}{4} \cdot 2^4 - 2 \cdot 2^3 + 2 \cdot 2^2 + 2\right) + \left(\frac{1}{4} \cdot 1^4 - 2 \cdot 1^3 + 2 \cdot 1^2 + 1\right)$
- ⁷ $\frac{9}{2}$

Alternative Method 3 for (b)

- ⁵ $\left| \int_1^2 \dots dx \right|$
- ⁶ $\left| \left(\frac{1}{4} \cdot 2^4 - 2 \cdot 2^3 + 2 \cdot 2^2 + 2\right) - \left(\frac{1}{4} \cdot 1^4 - 2 \cdot 1^3 + 2 \cdot 1^2 + 1\right) \right|$
- ⁷ $\frac{9}{2}$

7 Solve the equation $\sin x^\circ - \sin 2x^\circ = 0$ in the interval $0 \leq x \leq 360$.

4

Qu.	part	marks	Grade	Syllabus Code	Calculator class	Source
7		4	C	T10	NC	06/46

The primary method m/s is based on the following generic m/s. THIS GENERIC M/S MAY BE USED AS AN EQUIVALENCE GUIDE BUT ONLY WHERE A CANDIDATE DOES NOT USE THE PRIMARY METHOD OR ANY ALTERNATIVE METHOD SHOWN IN DETAIL IN THE MARKING SCHEME

- ¹ ss know to use double angle formula
- ² pr factorise
- ³ pr solve
- ⁴ ic know exact values

Primary Method : Give 1 mark for each •

- ¹ $\sin(x^\circ) - 2\sin(x^\circ)\cos(x^\circ) = 0$
- ² $\sin(x^\circ)(1 - 2\cos(x^\circ)) = 0$
- ³ $\sin(x^\circ) = 0$ or $\cos(x^\circ) = 0.5$
- ⁴ $x = 0, 180, 360, \quad 60, 300$

4

Notes

- 1 An "=" 0" must appear somewhere between the start and •² evidence.
- 2 The inclusion of extra answers which would have been correct with a larger interval should be treated as bad form and NOT penalised.
- 3 The omission of a correct answer (e.g. 0) means the candidates loses a mark (•⁴ in the Primary Method).
- 4 Candidates may embark on a journey with the wrong formula for $\sin(2x^\circ)$. With an equivalent level of difficulty it may still be worth a maximum of 3 marks. See Common Error No.1.
- 5 Candidates who draw a sketch of $y = \sin(x^\circ)$ and $y = \sin(2x^\circ)$ giving 0,180,360 may be awarded •¹ and •³.

Alternative Marking Method (Cross marking for •3 and •4)

- ¹ $\sin(x^\circ) - 2\sin(x^\circ)\cos(x^\circ) = 0$
- ² $\sin(x^\circ)(1 - 2\cos(x^\circ)) = 0$
- ³ $\sin(x^\circ) = 0$ and $x = 0, 180, 360$
- ⁴ $\cos(x^\circ) = 0.5$ and $x = 60, 300$

Alternative Method Division by $\sin(x)$

- ¹ $\sin(x^\circ) - 2\sin(x^\circ)\cos(x^\circ) = 0$
- ² either $\sin(x^\circ) = 0$ or $\sin(x^\circ) \neq 0$
- ³ $\sin(x^\circ) = 0 \Rightarrow x = 0, 180, 360$
- ⁴ $\cos(x^\circ) = 0.5 \Rightarrow x = 60, 300$

Common Error No.1

X •¹ $\sin(x^\circ) - (1 - 2\sin^2(x^\circ)) = 0$
 $2\sin^2(x^\circ) + \sin(x^\circ) - 1 = 0$
 X ✓ •² $(2\sin(x^\circ) - 1)(\sin(x^\circ) + 1) = 0$
 X ✓ •³ $\sin(x^\circ) = \frac{1}{2}$ or $\sin(x^\circ) = -1$
 X ✓ •⁴ $x = 30, 150, \quad x = 270$
 award 3 marks

Common Error No.2

$\sin(x^\circ) - \sin^2(x^\circ) = 0$
 X •¹
 X ✓ •² $\sin(x^\circ)(1 - \sin(x^\circ)) = 0$
 X •³ $\sin(x^\circ) = 0$ or $\sin(x^\circ) = 1$
 X ✓ •⁴ $x = 0, 180, 360, \quad 90$
 award 2 marks

Common Error No.3

$\sin(x) - \sin(2x) = 0$
 $\sin(x) = 0, \sin(2x) = 0$
 etc
 gains NO marks

- 8 (a) Express $2x^2 + 4x - 3$ in the form $a(x + b)^2 + c$. 3
- (b) Write down the coordinates of the turning point on the parabola with equation $y = 2x^2 + 4x - 3$. 1

Qu.	part	marks	Grade	Syllabus Code	Calculator class	Source
8	a	3	B	A5	NC	06/32
	b	1	C	A6	NC	

The primary method m/s is based on the following generic m/s. THIS GENERIC M/S MAY BE USED AS AN EQUIVALENCE GUIDE BUT ONLY WHERE A CANDIDATE DOES NOT USE THE PRIMARY METHOD OR ANY ALTERNATIVE METHOD SHOWN IN DETAIL IN THE MARKING SCHEME

- ¹ ss know how to complete (deal with the "a")
- ² pr process the value of "b"
- ³ pr process the value of "c"
- ⁴ ic interpret equation of parabola

Primary Method : Give 1 mark for each

- ¹ $a = 2$
- ² $b = 1$
- ³ $c = -5$ 3
- ⁴ $(-1, -5)$ 1

Note

- 1 Alternative Method 1 should be used for assessing part marks/follow throughs.
- 2 For •⁴, no justification is required. Candidates may choose to differentiate etc. but may still earn only one mark for the correct answer.
- 3 For •⁴, accept $(-b, c)$.


Alternative Method 1 for (a)

- ¹ $2(x^2 + 2x)$
- ² $2(x + 1)^2$
- ³ $2(x + 1)^2 - 5$
- ⁴ $(-1, -5)$

Alternative Method 2 for (a) : Comparing coefficients

- ¹ $2x^2 + 4x - 3 = ax^2 + 2abx + ab^2 + c \Rightarrow a = 2$
- ² $2ab = 4 \Rightarrow b = 1$
- ³ $ab^2 + c = -3 \Rightarrow c = -5$
- ⁴ $(-1, -5)$

9 u and v are vectors given by $u = \begin{pmatrix} k^3 \\ 1 \\ k+2 \end{pmatrix}$ and $v = \begin{pmatrix} 1 \\ 3k^2 \\ -1 \end{pmatrix}$, where $k > 0$.



- (a) If $u \cdot v = 1$ show that $k^3 + 3k^2 - k - 3 = 0$. 2 marks
- (b) Show that $(k + 3)$ is a factor of $k^3 + 3k^2 - k - 3$ and hence factorise $k^3 + 3k^2 - k - 3$ fully. 5 marks
- (c) Deduce the only possible value of k . 1 mark
- (d) The angle between u and v is θ . Find the exact value of $\cos \theta$. 3 marks

Qu.	part	marks	Grade	Syllabus Code	Calculator class	Source
8	a	2	C	G26	CN	05/10
	b	5	C	A21	NC	
	c	1	C	A6	CN	
	d	3	C	G28	NC	

The primary method m/s is based on the following generic m/s. THIS GENERIC M/S MAY BE USED AS AN EQUIVALENCE GUIDE BUT ONLY WHERE A CANDIDATE DOES NOT USE THE PRIMARY METHOD OR ANY ALTERNATIVE METHOD SHOWN IN DETAIL IN THE MARKING SCHEME

- ¹ pr find scalar product
- ² ic complete proof
- ³ ss know to use $k = -3$
- ⁴ pr complete evaluation and conclusion
- ⁵ ic start to find quadratic factor
- ⁶ ic complete quadratic factor
- ⁷ pr factorise completely
- ⁸ ic interpret k
- ⁹ ic interpret vectors
- ¹⁰ pr find magnitudes
- ¹¹ ss use formula

Notes

- 1 No explanation is required for k but the chosen value must follow from the working for •⁶ or •⁷. **Do not accept $\sqrt{1}$.**
- 2 In primary method (•⁴) and alternative (•⁵) candidates must show some acknowledgement of the resulting "zero". Although a statement w.r.t. the zero is preferable, accept something as simple as "underlining" the zero.
- 3 Only numerical values are acceptable for •⁹, •¹⁰ and •¹¹; answers are acceptable in unsimplified form eg $\cos \theta = \frac{1}{\sqrt{11} \times \sqrt{11}}$

Alternative method 1 (marks 3–7) Long Division

•³ $k+3 \overline{) \begin{matrix} k^3 & +3k^2 & -k & -3 \\ k^3 & +3k^2 & & \\ \hline & & -k & -3 \\ & & -k & -3 \\ \hline & & & 0 \end{matrix}}$

•⁴ $\underline{-k \quad -3}$

•⁵ remainder is zero so $(k+3)$ is a factor

•⁶ $k^2 - 1$

•⁷ $(k+3)(k+1)(k-1)$ **stated explicitly**

Primary Method : Give 1 mark for each

- ¹ $u \cdot v = k^3 \cdot 1 + 1 \cdot (3k^2) + (k+2) \cdot (-1)$ stated or implied by •² before completion
 - ² $k^3 + 3k^2 - k - 2 = 1$ and complete 2 marks
 - ³ know to use $k = -3$
 - ⁴ $-27 + 27 - (-3) - 3 = 0 \Rightarrow x+3$ is a factor
 - ⁵ $(k+3)(k^2 \dots)$
 - ⁶ $(k+3)(k^2 - 1)$
 - ⁷ $(k+3)(k+1)(k-1)$ **stated explicitly** 5 marks
 - ⁸ $k = 1$ 1 mark
 - ⁹ $u = \begin{pmatrix} 1 \\ 1 \\ 3 \end{pmatrix}, v = \begin{pmatrix} 1 \\ 3 \\ -1 \end{pmatrix}$ **stated or implied by •¹⁰**
 - ¹⁰ $|u| = \sqrt{11}$ and $|v| = \sqrt{11}$
 - ¹¹ $\cos \theta = \frac{1}{11}$ 3 marks
- N.B.**
•⁹ and •¹⁰ may be cross-marked.

Alternative method 2 (marks 3–7) Synthetic Division

•³ $\begin{array}{r|rrrr} -3 & & & & \\ \hline & & & & \end{array}$


•⁴ $\begin{array}{r|rrrr} -3 & 1 & 3 & -1 & -3 \\ \hline & & -3 & 0 & 3 \\ \hline & 1 & 0 & -1 & 0 \end{array}$

•⁵ " $f(-3) = 0$ " so $(k+3)$ is a factor

•⁶ $(k^2 - 1)$

•⁷ $(k+3)(k+1)(k-1)$ **stated explicitly**

- 10 Two variables, x and y , are connected by the law $y = a^x$. A graph of $\log_4(y)$ against x is a straight line passing through the origin and the point A(6,3). Find the value of a .



4

Qu.	part	marks	Grade	Syllabus Code	Calculator class	Source
10		4	A	A33	NC	06/91

The primary method m/s is based on the following generic m/s. THIS GENERIC M/S MAY BE USED AS AN EQUIVALENCE GUIDE BUT ONLY WHERE A CANDIDATE DOES NOT USE THE PRIMARY METHOD OR ANY ALTERNATIVE METHOD SHOWN IN DETAIL IN THE MARKING SCHEME

- ¹ ss know to take logarithms
- ² ic substitute known point
- ³ pr solve
- ⁴ pr solve

Primary Method : Give 1 mark for each •

- ¹ $\log_4(y) = \log_4(a^x)$
- ² $3 = \log_4(a^6)$
- ³ $a^6 = 4^3$
- ⁴ $a = 2$

4 marks

Note

- 1 $m = \frac{1}{2}$ and nothing else gains no marks.
- 2 For •⁴, a correct answer without any legitimate evidence gains **NO** marks.
- 3 For •⁴, ignore the inclusion of a negative answer.

Alternative Method 1

- ¹ $\log_4(y) = \log_4(a^x)$
- ² $3 = 6 \log_4(a)$
- ³ $\log_4(a) = \frac{1}{2}$
- ⁴ $a = 2$

Alternative Method 2

- ¹ $\log_4(y) = mx + c$
- ² $m = \frac{1}{2}, c = 0$
- ³ $y = 4^{\frac{1}{2}x}$
- ⁴ $y = \left(4^{\frac{1}{2}}\right)^x = 2^x \Rightarrow a = 2$

Common Error 1

- | | | |
|---|----------------|---------------------------|
| ✓ | • ¹ | $\log_4(y) = \log_4(a^x)$ |
| X | • ² | $\log_4(3) = \log_4(a^6)$ |
| X | • ³ | $3 = a^6$ |
| X | • ⁴ | $a = 3^{\frac{1}{6}}$ |

Alternative Method 3

- ¹ At A $\log_4(y) = 3$
- ² $y = 4^3$
- ³ $a^6 = 4^3$
- ⁴ $a = 2$

Common Error 2

- | | | |
|---|----------------|-----------------|
| X | • ¹ | $\log_4(y) = x$ |
| X | • ² | -- |
| X | • ³ | $y = 4^x$ |
| X | • ⁴ | $a = 4$ |

Alternative Method 4

- ¹ $\log_4(y) = \log_4(a^x)$
- ² $\log_4(y) = x \log_4(a)$
- ³ $\log_4(a) = \frac{1}{2}$
- ⁴ $a = 4^{\frac{1}{2}} = 2$