



2012 Mathematics

Advanced Higher

Finalised Marking Instructions

© Scottish Qualifications Authority 2012

The information in this publication may be reproduced to support SQA qualifications only on a non-commercial basis. If it is to be used for any other purposes written permission must be obtained from SQA's NQ Delivery: Exam Operations.

Where the publication includes materials from sources other than SQA (secondary copyright), this material should only be reproduced for the purposes of examination or assessment. If it needs to be reproduced for any other purpose it is the centre's responsibility to obtain the necessary copyright clearance. SQA's NQ Delivery: Exam Operations may be able to direct you to the secondary sources.


These Marking Instructions have been prepared by Examination Teams for use by SQA Appointed Markers when marking External Course Assessments. This publication must not be reproduced for commercial or trade purposes.

Advanced Higher Mathematics 2012

|Marks awarded for

1.	(3,4)	<p>(a) $f(x) = \frac{3x + 1}{x^2 + 1}$</p> $f'(x) = \frac{3(x^2 + 1) - (3x + 1)2x}{(x^2 + 1)^2}$ $= \frac{3x^2 + 3 - 6x^2 - 2x}{(x^2 + 1)^2}$ $= \frac{-3x^2 - 2x + 3}{(x^2 + 1)^2}$		<p>1M for quotient rule (or product)</p> <p>1 for two correct terms</p> <p>1 for third correct term</p>
	(b)	<p>$g(x) = \cos^2 x e^{\tan x}$</p> $g'(x) = 2 \cos x (-\sin x) e^{\tan x} + (\cos^2 x)(\sec^2 x) e^{\tan x}$ $= -\sin 2x e^{\tan x} + e^{\tan x}$ $= (1 - \sin 2x) e^{\tan x}$		<p>1M product rule</p> <p>1 first correct term</p> <p>1 second correct term</p> <p>1 simplification</p>
	(b) alternative	<p>$g(x) = \cos^2 x \exp(\tan x)$</p> $\ln(g(x)) = \ln(\cos^2 x) + \tan x$ $= 2 \ln(\cos x) + \tan x$		1M
		<p>Differentiating</p> $\frac{g'(x)}{g(x)} = 2 \frac{(-\sin x)}{\cos x} + \sec^2 x$		1
		$g'(x) = \left(\frac{1 - 2 \sin x \cos x}{\cos^2 x} \right) \cos^2 x \exp(\tan x)$ $= (1 - \sin 2x) \tan x$		1

2.	(5)	<p>$a = 2048$ and $ar^3 = 256$</p> $\Rightarrow r^3 = \frac{1}{8}$ $\Rightarrow r = \frac{1}{2}$ $S_n = \frac{a(1 - r^n)}{1 - r}$ $\Rightarrow \frac{1 - (\frac{1}{2})^n}{1 - \frac{1}{2}} = \frac{4088}{2048}$ $= \frac{511}{256}$ $\Rightarrow 1 - \left(\frac{1}{2}\right)^n = \frac{511}{256} \times \frac{1}{2} = \frac{511}{512}$ $\frac{1}{2^n} = 1 - \frac{511}{512} = \frac{1}{512}$ $\Rightarrow 2^n = 512 \Rightarrow n = 9$		<p>1M valid approach</p> <p>1 correct answer only, 2 marks</p> <p>1M for sum formula</p> <p>1</p> <p>1 any valid method</p>
-----------	------------	---	--	--

<p>3. Since w is a root, $\bar{w} = -1 - 2i$ is also a root.</p> <p>(6) The corresponding factors are $(z + 1 - 2i)$ and $(z + 1 + 2i)$ from which $((z + 1) - 2i)((z + 1) + 2i) = (z + 1)^2 + 4$ $= z^2 + 2z + 5$ $z^3 + 5z^2 + 11z + 15 = (z^2 + 2z + 5)(z + 3)$</p> <p>The roots are $(-1 + 2i)$, $(-1 - 2i)$ and -3.</p> 	<p>1 for conjugate</p> <p>1 evidence needed</p> <p>1 for stating roots together</p> <p>1 for two correct points</p> <p>1 for third correct point</p>
---	---

<p>4. The general term is given by:</p> <p>(5)</p> $\binom{9}{r} (2x)^{9-r} \left(-\frac{1}{x^2}\right)^r$ $= \binom{9}{r} \times \frac{2^{9-r} x^{9-r} (-1)^r}{x^{2r}}$ $= \binom{9}{r} \times (-1)^r 2^{9-r} x^{9-3r}$ <p>The term independent of x occurs when $9 - 3r = 0$, i.e. when $r = 3$.</p> <p>The term is: $\frac{9!}{6! 3!} (-1)^3 2^6$ $= -5376$.</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
--	---

<p>5. Method 1</p> <p>(5) $\vec{PQ} = 3\mathbf{i} + \mathbf{j} + 4\mathbf{k}$ and $\vec{QR} = 2\mathbf{i} - 2\mathbf{j} - 2\mathbf{k}$ A normal to the plane:</p> $\vec{PQ} \times \vec{QR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & 1 & 4 \\ 2 & -2 & -2 \end{vmatrix}$ $= \mathbf{i} \begin{vmatrix} 1 & 4 \\ -2 & -2 \end{vmatrix} - \mathbf{j} \begin{vmatrix} 3 & 4 \\ 2 & -2 \end{vmatrix} + \mathbf{k} \begin{vmatrix} 3 & 1 \\ 2 & -2 \end{vmatrix}$ $= 6\mathbf{i} + 14\mathbf{j} - 8\mathbf{k}$ <p>Hence the equation has the form: $6x + 14y - 8z = d$.</p> <p>The plane passes through $P(-2, 1, -1)$ so $d = -12 + 14 + 8 = 10$ which gives an equation $6x + 14y - 8z = 10$ i.e. $3x + 7y - 4z = 5$.</p>	<p>1 \vec{PR} could be used</p> <p>1M</p> <p>1</p> <p>1</p> <p>1</p>
---	--

Method 2

A plane has an equation of the form

 $ax + by + cz = d$. Using the points P, Q, R we get

$$-2a + b - c = d$$

$$a + 2b + 3c = d$$

$$3a + c = d$$

1M

Using Gaussian elimination to solve these we have

$$\begin{vmatrix} -2 & 1 & -1 & d \\ 1 & 2 & 3 & d \\ 3 & 0 & 1 & d \end{vmatrix} \Rightarrow \begin{vmatrix} -2 & 1 & -1 & d \\ 0 & 5 & 5 & 3d \\ 0 & 6 & 8 & 2d \end{vmatrix} \quad \mathbf{1}$$

$$\Rightarrow \begin{vmatrix} -2 & 1 & -1 & d \\ 0 & 5 & 5 & 3d \\ 0 & 0 & 2 & -\frac{8}{5}d \end{vmatrix} \quad \mathbf{1}$$

$$\Rightarrow c = -\frac{4}{5}d, \quad b = \frac{7}{5}d, \quad a = \frac{3}{5}d \quad \mathbf{1}$$

These give the equation

$$\left(\frac{3}{5}d\right)x + \left(\frac{7}{5}d\right)y + \left(-\frac{4}{5}d\right)z = d$$

i.e. $3x + 7y - 4z = 5$

1

or other valid method

6. Method 1

(5) $e^x = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \dots$ **1**
 $(1 + e^x)^2 = 1 + 2e^x + e^{2x}$ **1M**
 $= 1 + 2\left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \dots\right)$ **1**
 $+ \left(1 + 2x + \frac{(2x)^2}{2} + \frac{(2x)^3}{6} + \dots\right)$ **1**
 $= 1 + 2 + 2x + x^2 + \frac{1}{3}x^3 + 1 + 2x + 2x^2 + \frac{4}{3}x^3 + \dots$ **1**
 $= 4 + 4x + 3x^2 + \frac{5}{3}x^3 + \dots$ **1**

Method 2


$e^x = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \dots$ **1**
 $(1 + e^x) = 2 + x + \frac{x^2}{2} + \frac{x^3}{6} + \dots$ **1**
 $(1 + e^x)^2 = (2 + x + \frac{x^2}{2} + \frac{x^3}{6} + \dots)(2 + x + \frac{x^2}{2} + \frac{x^3}{6} + \dots)$ **1M**
 $= 4 + 4x + 3x^2 + \frac{1}{3}x^3 + \frac{1}{2}x^3 + \frac{1}{2}x^3 + \frac{1}{3}x^3 + \dots$ **1**
 $= 4 + 4x + 3x^2 + \frac{5}{3}x^3 + \dots$ **1**

Method 3


$e^x = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \dots$ **1**
 $f(x) = (1 + e^x)^2$ $f(0) = 4$
 $f'(x) = 2e^x(1 + e^x)$ $f'(0) = 4$ **1**
 $= 2e^x + 2e^{2x}$
 $f''(x) = 2e^x + 4e^{2x}$ $f''(0) = 6$ **1**
 $f'''(x) = 2e^x + 8e^{2x}$ $f'''(0) = 10$ **1**
 $f(x) = f(0) + f'(0)x + \frac{f''(0)}{2}x^2 + \frac{f'''(0)}{6}x^3 + \dots$
 $(1 + e^x)^2 = 4 + 4x + 3x^2 + \frac{5}{3}x^3 + \dots$ **1**

can award marks for correct columns.

7. (4)



1 for shape
1 for coordinates



1 for both horizontal lines
1 for values: 1, -1, -2

8. (6)	$x = 4 \sin \theta \Rightarrow dx = 4 \cos \theta d\theta$	1	
	$x = 0 \Rightarrow \theta = 0$	1	
	$x = 2 \Rightarrow \theta = \frac{\pi}{6}$		
	$\int_0^2 \sqrt{16-x^2} dx$		
	$= \int_0^{\pi/6} \sqrt{16-(4 \sin \theta)^2} \cdot 4 \cos \theta d\theta$	1	
	$= \int_0^{\pi/6} \sqrt{16(1-\sin^2 \theta)} \cdot 4 \cos \theta d\theta$		
	$= \int_0^{\pi/6} \sqrt{16 \cos^2 \theta} \cdot 4 \cos \theta d\theta$		
	$= \int_0^{\pi/6} 16 \cos^2 \theta d\theta$	1	
	$= 8 \int_0^{\pi/6} (1 + \cos 2\theta) d\theta$		
	$= 8 [\theta + \frac{1}{2} \sin 2\theta]_0^{\pi/6}$	1	for applying trig. identity and integrating
$= \frac{8\pi}{6} + 4 \sin \frac{\pi}{3}$			
$= \frac{4\pi}{3} + 2\sqrt{3} (\approx 7.65)$	1	numerical approx. allowed	

9. (4)	<i>Method 1</i>		
	$A + A^{-1} = I$		
	$A^2 + I = A$	1	for multiplying by A
	Hence $A^2 + I = I - A^{-1}$	1	for rearranging $A + A^{-1} = I$
	$A^2 = -A^{-1}$	1	for subtracting I
	$A^3 = -I, \text{ i.e. } k = -1$	1	for multiplying by A
	<i>Method 2</i>		
	$A + A^{-1} = I$		
	$A = I - A^{-1}$	1	for rearranging
	$A^2 = I - 2A^{-1} + (A^{-1})^2$	1	for squaring
$A^3 = A - 2I + A^{-1}$	1	for multiplying by A	
$A^3 = (A + A^{-1}) - 2I = I - 2I$			
Hence $A^3 = -I, \text{ i.e. } k = -1$	1		
<i>Method 3</i>			
$A + A^{-1} = I$			
$A = I - A^{-1}$	1	for rearranging	
$A^3 = A^2 - A$	1	for multiplying by A^2	
$A^3 = (A - I) - A$	1	using $A^2 = A - I$	
$= -I, \text{ i.e. } k = -1$	1		
<i>Plus other valid methods.</i>			

10.	<i>Method 1</i>	$1234 = 7 \times 176 + 2$	1	answer only, 1 of 3
(3)		$176 = 7 \times 25 + 1$	1	
		$25 = 7 \times 3 + 4$	1	
	Hence	$1234_{10} = 3412_7$	1	
	<i>Method 2</i>			
		$1234 = 7 \times 176 + 2$	1	
		$= 7 \times (7 \times 25 + 1) + 2$		
		$= 7 \times (7 \times (7 \times 3 + 4) + 1) + 2$	1	
		$= 3 \times 7^3 + 4 \times 7^2 + 1 \times 7 + 2$		
	Hence	$1234_{10} = 3412_7$	1	answer only, 1 of 3
11.	(a)	$\frac{d}{dx}(\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}$	1	
(1,4)	(b) $\int \sin^{-1} x \cdot \frac{x}{\sqrt{1-x^2}} dx =$			
		$\sin^{-1} x \int \frac{x}{\sqrt{1-x^2}} dx - \int \left(\frac{d}{dx}(\sin^{-1} x) \int \frac{x}{\sqrt{1-x^2}} dx \right) dx$	1	
		$= \sin^{-1} x \int \frac{x}{\sqrt{1-x^2}} dx - \int \left(\frac{1}{\sqrt{1-x^2}} \int \frac{x}{\sqrt{1-x^2}} dx \right) dx$		
		$= \sin^{-1} x (-\sqrt{1-x^2}) - \int \left(\frac{1}{\sqrt{1-x^2}} (-\sqrt{1-x^2}) \right) dx$	1	for $\int \frac{x}{\sqrt{1-x^2}} dx = -\sqrt{1-x^2}$
		$= \sin^{-1} x (-\sqrt{1-x^2}) - \int (-1) dx$	1	
		$= x - \sin^{-1} x \sqrt{1-x^2} + c$	1	
12.	(5)	$\frac{dr}{dt} = -0.02; \quad \frac{dh}{dt} = 0.01$	1	
		$V = \pi r^2 h$		
		$\frac{dV}{dt} = \pi \left(2r \frac{dr}{dt} \right) h + \pi r^2 \frac{dh}{dt}$	1M	for implicit differentiation
		$= \pi (2 \times 0.6 \times (-0.02) \times 2 + 0.36 \times 0.01)$	1	for accuracy
		$= \pi (-0.048 + 0.0036)$		
		$= -0.0444\pi (\approx -0.14)$		
	The rate of change in the volume is	$-0.0444\pi \text{ m}^3 \text{ s}^{-1}$.	1	units required

13.
(10)

$$x = 2t + \frac{1}{2}t^2 \quad \Rightarrow \quad \frac{dx}{dt} = 2 + t \quad \mathbf{1}$$

$$y = \frac{1}{3}t^3 - 3t \quad \Rightarrow \quad \frac{dy}{dt} = t^2 - 3 \quad \mathbf{1}$$

$$\frac{dy}{dx} = \frac{t^2 - 3}{2 + t} \quad \mathbf{1}$$

$$\frac{d}{dt} \left(\frac{dy}{dx} \right) = \frac{2t(2+t) - (t^2-3)}{(2+t)^2} = \frac{t^2 + 4t + 3}{(2+t)^2} \quad \mathbf{1}$$

$$\frac{d^2y}{dx^2} = \frac{t^2 + 4t + 3}{(2+t)^2} \times \frac{1}{2+t} = \frac{t^2 + 4t + 3}{(2+t)^3} \quad \mathbf{1}$$

Stationary points when $\frac{dy}{dx} = 0$, i.e.

$$t^2 - 3 = 0 \Rightarrow t = \pm\sqrt{3} \quad \mathbf{1}$$

When $t = \sqrt{3}$, $\frac{d^2y}{dx^2} = \frac{3 + 4\sqrt{3} + 3}{(2 + \sqrt{3})^3} > 0$

which gives a minimum. **1**

When $t = -\sqrt{3}$, $\frac{d^2y}{dx^2} = \frac{3 - 4\sqrt{3} + 3}{(2 - \sqrt{3})^3} < 0$

which gives a maximum. **1**

At a point of inflexion, $\frac{d^2y}{dx^2} = 0$. **1**

In this case, that means

$$t^2 + 4t + 3 = (t + 1)(t + 3) = 0$$

and this has exactly two roots. **1**

Note that this is a slimmed-down version of the complete story of points of inflexion.

no marks for using a nature table

no marks for using a nature table

need to show 2 values exist

14.
(5,
1,3)

(a)

$$\left| \begin{array}{ccc|c} 4 & 0 & 6 & 1 \\ 2 & -2 & 4 & -1 \\ -1 & 1 & \lambda & 2 \end{array} \right|$$

1 for augmented matrix

$$\left| \begin{array}{ccc|c} 4 & 0 & 6 & 1 \\ 0 & 4 & -2 & 3 \\ 0 & 4 & 6+4\lambda & 9 \end{array} \right|$$

1

$$\left| \begin{array}{ccc|c} 4 & 0 & 6 & 1 \\ 0 & 4 & -2 & 3 \\ 0 & 0 & 8+4\lambda & 6 \end{array} \right|$$

1 triangular form needed

$$z = \frac{6}{8+4\lambda} = \frac{3}{2(2+\lambda)}$$

1 first root

$$4y = 3 + 2z \Rightarrow 4y = \frac{18+6\lambda}{4+2\lambda}$$

$$\Rightarrow y = \frac{3\lambda+9}{4(2+\lambda)}$$

$$4x = 1 - 6z \Rightarrow 4x = \frac{2\lambda-14}{4+2\lambda}$$

$$\Rightarrow x = \frac{\lambda-7}{4(2+\lambda)}$$

1 other two roots

(b) When $\lambda = -2$, the final row gives $0 = 6$ which is inconsistent.

There are no solutions.

1

(c) $\lambda = -2.1$; $x = 22.75$; $y = -6.75$; $z = -15$

1,1 1 for first 2 values; 1 for third

Although the values of λ are close, the values of x , y and z are quite different. The

system is **ill-conditioned** near $\lambda = -2$.

1

15.
(4,7) (a) $\frac{1}{(x-1)(x+2)^2} = \frac{A}{x-1} + \frac{B}{x+2} + \frac{C}{(x+2)^2}$ **1M**

$$1 = A(x+2)^2 + B(x-1)(x+2) + C(x-1)$$

$$x = 1 \Rightarrow A = \frac{1}{9} \quad \mathbf{1}$$

$$x = -2 \Rightarrow C = -\frac{1}{3} \quad \mathbf{1}$$

$$x = 0 \Rightarrow 1 = \frac{4}{9} - 2B + \frac{1}{3} \Rightarrow B = -\frac{1}{9} \quad \mathbf{1}$$

$$\therefore \frac{1}{(x-1)(x+2)^2} = \frac{1}{9} \left(\frac{1}{x-1} - \frac{1}{x+2} - \frac{3}{(x+2)^2} \right)$$

(b) $(x-1) \frac{dy}{dx} - y = \frac{x-1}{(x+2)^2}$

$$\frac{dy}{dx} - \frac{1}{x-1}y = \frac{1}{(x+2)^2} \quad \mathbf{1M} \text{ for rearranging}$$

Integrating factor: $\exp\left(\int -\frac{1}{x-1}dx\right)$ **1**

$$= \exp(-\ln(x-1)) = (x-1)^{-1} \quad \mathbf{1}$$

$$\frac{1}{(x-1)} \frac{dy}{dx} - \frac{1}{(x-1)^2}y = \frac{1}{(x-1)(x+2)^2}$$

$$\frac{d}{dx} \left(\frac{y}{x-1} \right) = \frac{1}{(x-1)(x+2)^2} \quad \mathbf{1}$$

$$= \frac{1}{9} \left(\frac{1}{x-1} - \frac{1}{x+2} - \frac{3}{(x+2)^2} \right) \quad \mathbf{1}$$

$$\frac{y}{x-1} = \frac{1}{9} \left(\ln|x-1| - \ln|x+2| + \frac{3}{x+2} \right) + c \quad \mathbf{1} \text{ constant of integration needed.}$$

$$y = \frac{x-1}{9} \left(\ln|x-1| - \ln|x+2| + \frac{3}{x+2} \right) + c(x-1) \quad \mathbf{1}$$

$$= \frac{x-1}{9} \left(\ln \left| \frac{x-1}{x+2} \right| + \frac{3}{x+2} \right) + c(x-1)$$

16. (6,4)	(a) For $n = 1$, the LHS = $\cos \theta + i \sin \theta$ and the RHS = $\cos \theta + i \sin \theta$. Hence the result is true for $n = 1$.	1	
	Assume the result is true for $n = k$, i.e. $(\cos \theta + i \sin \theta)^k = \cos k\theta + i \sin k\theta$.	1	working with n is penalised.
	Now consider the case when $n = k + 1$: $(\cos \theta + i \sin \theta)^{k+1} = (\cos \theta + i \sin \theta)^k (\cos \theta + i \sin \theta)$ $= (\cos k\theta + i \sin k\theta)(\cos \theta + i \sin \theta)$	1 1	for applying the inductive hypothesis
	$= (\cos k\theta \cos \theta - \sin k\theta \sin \theta) + i(\sin k\theta \cos \theta + \cos k\theta \sin \theta)$ $= \cos(k+1)\theta + i \sin(k+1)\theta$	1	multiplying and collecting
	Thus, if the result is true for $n = k$ the result is true for $n = k + 1$.		
	Since it is true for $n = 1$, the result is true for all $n \geq 1$.	1	
	(b) $\frac{(\cos \frac{\pi}{18} + i \sin \frac{\pi}{18})^{11}}{(\cos \frac{\pi}{36} + i \sin \frac{\pi}{36})^4} = \frac{\cos \frac{11\pi}{18} + i \sin \frac{11\pi}{18}}{\cos \frac{\pi}{9} + i \sin \frac{\pi}{9}}$ $= \frac{\cos \frac{11\pi}{18} + i \sin \frac{11\pi}{18}}{\cos \frac{\pi}{9} + i \sin \frac{\pi}{9}} \times \frac{\cos \frac{\pi}{9} - i \sin \frac{\pi}{9}}{\cos \frac{\pi}{9} - i \sin \frac{\pi}{9}}$ $= \frac{\cos \frac{11\pi}{18} \cos \frac{\pi}{9} + \sin \frac{11\pi}{18} \sin \frac{\pi}{9}}{\cos^2 \frac{\pi}{9} + \sin^2 \frac{\pi}{9}} + \text{imaginary term}$ $= \cos\left(\frac{11\pi}{18} - \frac{\pi}{9}\right) + \text{imaginary term}$ $= \cos \frac{\pi}{2} + \text{imaginary term}$	1 1 1	using result from above
	Thus the real part is zero as required.	1	or equivalent

END OF SOLUTIONS