

## Quadratic Functions

### Solving Quadratic Equations (Revision from Nat 5)

Quadratic equations must be solved by factorising when one side is equal to zero.

**Example 1:** Solve:

a)  $x^2 - 5x + 4 = 0$

b)  $3x^2 - 4x - 4 = 0$

c)  $(x - 5)(x + 2) = 4$

d)  $4x^2 - 3x - 2 = 0$


(answer to 2 d.p.)

### Finding the Equation of a Quadratic Function From Its Graph: $y = k(x - a)(x - b)$


If the graph of a quadratic function has roots at  $x = -1$  and  $x = 5$ , a reasonable guess at its equation would be  $y = x^2 - 4x - 5$ , i.e. from  $y = (x + 1)(x - 5)$ .

However, as the diagram shows, there are many parabolas which pass through these points, all of which belong to the **family** of functions  $y = k(x + 1)(x - 5)$ .

To find the equation of the original function, we need the roots and one other point on the curve (to allow us to determine the value of  $k$ ).



**Example 2:** State the equation of the graph below in the form  $y = ax^2 + bx + c$ .




## Completing the Square (Revision)

The diagram shows the graphs of two quadratic functions.

If the graph of  $y = x^2$  is shifted  $q$  units to the right, followed by  $r$  units up, then the graph of  $y = (x - q)^2 + r$  is obtained.

As the turning point of  $y = x^2$  is  $(0, 0)$ , it follows that the new curve has a turning point at  $(q, r)$ .

A quadratic equation written as  $y = p(x - q)^2 + r$  is said to be in the **completed square form**.



**Example 3:** (i) Write the following in the form  $y = (x + q)^2 + r$  and find the coordinates of the TP.

(ii) Hence state the minimum value of  $y$  and the corresponding value of  $x$ .

a)  $y = x^2 + 6x + 10$

b)  $y = x^2 - 3x + 1$

## Completing the Square when the $x^2$ Coefficient $\neq 1$

**Example 4:** Write  $y = 3x^2 + 12x + 5$  in the form  $y = p(x + q)^2 + r$ .

**Example 5:** Write  $y = 5 + 12x - x^2$  in the form  $y = p - (x + q)^2$ .

**Example 6:**

a) Write  $y = x^2 - 10x + 28$  in the form  $y = (x + p)^2 + q$ .


b) Hence find the maximum value of  $\frac{18}{x^2 - 10x + 28}$

## Solving Quadratic Inequations

Quadratic inequations are easily solved by making a sketch of the equivalent quadratic function, then determining the regions above or below the  $x$  - axis.

**Example 7:** Find the values of  $x$  for which: a)  $2x^2 - 7x + 6 > 0$  b)  $2x^2 - 7x + 6 < 0$

First, sketch  $y = 2x^2 - 7x + 6$



## Roots of Quadratic Equations and The Discriminant (Revision)

For  $y = ax^2 + bx + c$ ,  $b^2 - 4ac$  is known as the discriminant.

- $b^2 - 4ac > 0$  gives real, unequal roots
- $b^2 - 4ac = 0$  gives real, equal roots
- $b^2 - 4ac < 0$  gives NO real roots

If  $b^2 - 4ac$  gives a perfect square, the roots are RATIONAL  
If  $b^2 - 4ac$  does NOT give a perfect square, the roots are IRRATIONAL (i.e. surds)

**Example 8:** Find the value(s) of  $p$  given that  $2x^2 + 4x + p = 0$  has real roots.




**Example 9:** Show that the roots of  $x^2 + px = 3 - x$  are always real for all values of  $p$ .

**Example 10:** Find the value(s) of  $r$  given that  $x^2 + (r - 3)x + r = 0$  has no real roots.

## Tangents to Curves: Using the Discriminant

To find the points of contact between a line and a curve, we make the curve and line equations equal (i.e. make  $y = y$ ) to obtain a quadratic equation, and solve to find the  $x$ -coordinates.

By finding the discriminant of this quadratic equation, we can work out **how many** points of contact there are between the line and the curve. There are 3 options:

		
<b>Two points of contact</b> 2 distinct roots $b^2 - 4ac > 0$	<b>One point of contact</b> Equal roots $b^2 - 4ac = 0$	<b>No points of contact</b> No real roots $b^2 - 4ac < 0$

The most common use for this technique is to show that a line is a tangent to a curve

**Example 11:** Find the value of  $c$  such that the line  $y = 3x - 13$  is a tangent to the curve  $y = x^2 - 7x + c$ , and state the coordinates of the point of contact.

**Example 12:** Find two values of  $m$  such that  $y = mx - 7$  is a tangent to  $y = x^2 + 2x - 3$

**Past Paper Example 1:** Express  $2x^2 + 12x + 1$  in the form  $a(x + b)^2 + c$ .

**Past Paper Example 2:** Given that  $2x^2 + px + p + 6 = 0$  has no real roots, find the range of values for  $p$ .

**Past Paper Example 3:** Show that the roots of  $(k - 2)x^2 - 3kx + 2k = -2x$  are always real.