


## The Circle

If we draw, suitable to relative axes, a circle, radius  $r$ , centred on the origin, then the distance from the centre of any point  $P(x, y)$  could be determined to be  $d = \sqrt{x^2 + y^2}$ .



As the shape is a circle, then this distance is equal to the radius. It therefore follows that:

Since  $r = \sqrt{x^2 + y^2}$ , then  $r^2 = x^2 + y^2$

Therefore,

The equation  $x^2 + y^2 = r^2$  describes a circle with centre  $(0, 0)$  and radius  $r$

**Example 1:** Write down the centre and radius of each circle.

a)  $x^2 + y^2 = 64$

b)  $x^2 + y^2 = 361$

c)  $x^2 + y^2 = \frac{3}{25}$


**Example 2:** State where the points  $(-2, 7)$ ,  $(6, -8)$  and  $(5, 9)$  lie in relation to the circle  $x^2 + y^2 = 100$ .

## Circles with Centres Not at the Origin

The radius in the above circle is the distance between  $(x, y)$  and the origin, i.e.  $r = \sqrt{(x-0)^2 + (y-0)^2}$ . If we move the centre to the point  $(a, b)$ , then  $r = \sqrt{(x-a)^2 + (y-b)^2}$ .

Squaring both sides, we can now also say that:

The equation  $(x - a)^2 + (y - b)^2 = r^2$  describes a circle with centre  $(a, b)$  and radius  $r$



**Example 3:** Write down the centre and radius of each circle.

a)  $(x - 1)^2 + (y + 3)^2 = 4$

b)  $(x + 9)^2 + (y - 2)^2 = 20$

a)  $(x - 5)^2 + y^2 = 400$

**Example 4:** A is the point (4, 9) and B is the point (-2, 1).

Find the equation of the circle for which AB is the diameter.

**Example 5:** Points P, Q and R have coordinates (-10, 2), (5, 7) and (6, 4) respectively.

a) Show that triangle PQR is right angled at Q.

b) Hence find the equation of the circle passing through points P, Q and R.

### The General Equation of a Circle

For the circle described in Example 3a, we could expand the brackets and simplify to obtain the equation  $x^2 + y^2 - 2x + 6y + 6 = 0$ , which would also describe a circle with centre (1, -3) and radius 2.

$$\text{For } x^2 + y^2 + 2gx + 2fy + c = 0,$$

$$(x^2 + 2gx) + (y^2 + 2fy) = -c$$

$$(x^2 + 2gx + g^2) + (y^2 + 2fy + f^2) = g^2 + f^2 - c$$

$$(x + g)^2 + (y + f)^2 = (g^2 + f^2 - c)$$

**Example 6:** Find the centre and radius of the circle with equation  $x^2 + y^2 - 4x + 8y - 5 = 0$

Therefore, the circle described by

$$x^2 + y^2 + 2gx + 2fy + c = 0$$

has centre  $(-g, -f)$  and  $r = \sqrt{g^2 + f^2 - c}$

**Example 7:** State the range of values of  $c$  such that the equation  $x^2 + y^2 - 4x + 6y + c = 0$  describes a circle.




**Example 8:** Circles A and B are congruent. Circle A has the equation  $x^2 + y^2 + 8x - 12y + 7 = 0$ .

Find the equation of Circle B given that A and B touch only at the point (2, 9).

## Intersection of Lines and Circles

As with parabolas, there are **three** possibilities when a line and a circle come into contact, and we can examine the roots of a rearranged quadratic equation to determine which has occurred. However:

**We CANNOT make the circle and line equations equal to each other: the line equation must be substituted INTO the circle equation to obtain our quadratic equation!**


		
<b>Two points of contact</b> <b>2 distinct roots</b> $b^2 - 4ac > 0$	<b>One point of contact</b> <b>Equal roots</b> $b^2 - 4ac = 0$	<b>No points of contact</b> <b>No real roots</b> $b^2 - 4ac < 0$

**As with parabolas, the most common use of this technique is to show tangency.**

**Example 9:** Find the coordinates of the points of intersection of the line  $y = 2x - 1$  and the circle  $x^2 + y^2 - 2x - 12y + 27 = 0$ .

**Example 10:** Show that the line  $y = 3x + 10$  is a tangent to the circle  $x^2 + y^2 - 8x - 4y - 20 = 0$  and establish the coordinates of the point of contact.

**Example 11:** Find the equations of the tangents to the circle  $x^2 + y^2 = 9$  from the point  $(0, 5)$ .




### Tangents to Circles at Given Points

**Remember:** at the point of contact, the radius and tangent meet at  $90^\circ$  (i.e., they are perpendicular).

To find a tangent at a given point:

- Find the centre of the circle
- Find the gradient of the radius (joining C and the given point)
- Find the gradient of the tangent (flip and make negative)
- Sub the gradient and the original point into  $y - b = m(x - a)$




**Example 12:** Find the equation of the tangent to  $x^2 + y^2 - 14x + 6y - 87 = 0$  at the point  $(-2, 5)$ .

**Past Paper Example 1:** A circle has centre C  $(-2, 3)$  and passes through point P  $(1, 6)$ .

a) Find the equation of the circle.

b) PQ is a diameter of the circle. Find the equation of the tangent to this circle at Q.



**Past Paper Example 2:**

a) Show that the line with equation  $y = 3 - x$  is a tangent to the circle with equation

$$x^2 + y^2 + 14x + 4y - 19 = 0$$


and state the coordinates of P, the point of contact.

b) Relative to a suitable set of coordinate axes, the diagram opposite shows the circle from a) and a second smaller circle with centre C.

The line  $y = 3 - x$  is a common tangent at the point P.

The radius of the larger circle is three times the radius of the smaller circle.

Find the equation of the smaller circle.



**Past Paper Example 3:** Given that the equation

$$x^2 + y^2 - 2px - 4py + 3p + 2 = 0$$

represents a circle, determine the range of values of  $p$ .