

Intermediate 2 Units 1, 2, 3 Paper 1 2003



Created by
Graduate Bsc (Hons) MathsSci (Open) GIMA

1. Given $(2a - b)(3a + 2b)$

Multiplying out and gathering like terms we have:

$$\begin{aligned} &(2a - b)(3a + 2b) \\ &= 2a(3a + 2b) - b(3a + 2b) \\ &= 6a^2 + 4ab - 3ab - 2b^2 \\ &= 6a^2 + ab - 2b^2 \end{aligned}$$

2. Given the two spinners.



- (a) Completing table we get:

	1	2	3	4	5
Red	R,1	R,2	R,3	R,4	R,5
Yellow	Y,1	Y,2	Y,3	Y,4	Y,5
Blue	B,1	B,2	B,3	B,4	B,5
Green	G,1	G,2	G,3	G,4	G,5

- (b) Probability that we have $p(\text{Red, Even}) = \frac{2}{20} = \frac{1}{10}$

Intermediate 2 Units 1, 2, 3 Paper 1 2003

Created by


Graduate Bsc (Hons) MathsSci (Open) GIMA

3. Given the diagram. The volume for this cone will be:

$$V = \frac{1}{3}\pi r^2 h$$

$$V = \frac{1}{3}\pi \times 10^2 \times 12$$

$$V = 1256\text{cm}^3$$



4. Given the stem leaf diagram represents waiting times for Quickcars:

Waiting times (minutes)

0	6 7
1	2 3 4
2	5 6 9 9
3	2 5 7
4	2 4

$$n = 14$$

1|3 represents 13 minutes

- (a) Calculating the median, lower and upper quartiles we have:

$$\text{median} = \frac{26 + 29}{2} = 27.5 \quad \text{lower} = 13 \quad \text{upper} = 35$$

- (b) Semi-interquartile range is: $\frac{y_2 - y_1}{x_2 - x_1} = \frac{Q_3 - Q_1}{2} = \frac{35 - 13}{2} = 11$


- (c) If FastCabs have a semi-interquartile of 2.5 then they are more consistent because data is less spread out.

Intermediate 2 Units 1, 2, 3 Paper 1 2003

Created by

Graduate Bsc (Hons) MathsSci (Open) GIMA

5. Given that the graph represents a function of the form $y = a \sin bx^\circ$



The values for a and b are 3 and 2 respectively.

6. Expressing $\frac{\sqrt{40}}{\sqrt{2}}$ as a surd in its simplest form we get:

$$\frac{\sqrt{40}}{\sqrt{2}} = \frac{\sqrt{4}\sqrt{10}}{\sqrt{2}} = \frac{2\sqrt{2}\sqrt{5}}{\sqrt{2}} = 2\sqrt{5}$$

- (b) Simplifying $\frac{2x+2}{(x+1)^2}$ we get:

$$\frac{2x+2}{(x+1)^2} = \frac{2\cancel{(x+1)}^1}{(\cancel{x+1})^2_1} = \frac{2}{x+1}$$

Intermediate 2 Units 1, 2, 3 Paper 1 2003

Created by
Graduate Bsc (Hons) MathsSci (Open) GIMA

7. Given the two concentric circles.

AB is a tangent to the small circle and a chord to the big circle.

Also AB measures 16cm.

Red values have been added to diagram since they are easily calculated.


Since AB is a tangent radius of the small circle makes a right angle with the chord AB as shown. It also bisects (halves) AB.

Using Pythagoras Theorem or recognising a Pythagorean triple we have:

$$c^2 = a^2 + b^2$$

$$c = \sqrt{8^2 + 6^2}$$

$$c = 10\text{cm}$$



Intermediate 2 Units 1, 2, 3 Paper 1 2003

Created by
Graduate Bsc (Hons) MathsSci (Open) GIMA

8. (a) Factorising $7 + 6x - x^2$
Using FOIL (or any other suitable method) we get:

$$\begin{aligned} 7 + 6x - x^2 \\ = (7 - x)(1 + x) \end{aligned}$$

- (b) The roots of $7 + 6x - x^2$ are:

$$\begin{aligned} (7 - x) = 0 & \quad (1 + x) = 0 \\ x = 7 & \quad x = -1 \end{aligned}$$

- (c) Given the graph of $7 + 6x - x^2$

Red values have been added to diagram using information obtained in part (b).

By symmetry the x coordinate is 3.

y coordinate is given by:

$$y = 7 + 6 \times 3 - 3^2 = 16$$

Maximum turning point occurs at (3, 16)

